

# Protocol 102A

by Charlotte Lackney

**Protocol 102A** is similar to Protocol 101 (CDS Protocol-C) except for the addition of unactivated 22.4% sodium chlorite solution (SCS / MMS) for ingestion purposes only. MMS should not normally be added to CDS when used outside the stomach. AKA **CDM** (CDS + MMS).

The main reason to add MMS to 3000 ppm CDS is because when CDS dosing gets too high, often mouth and throat irritation will happen and the user can not increase dosing any higher. Another reason to add MMS to CDS for ingestion, even in stomachs with below normal gastric acids, is that MMS may be activated elsewhere in the body. (Needed for Autism recovery)

As you may know, each milliliter of 3000 ppm CDS contains 3 mg of chlorine dioxide (CLO<sub>2</sub>) and each drop of MMS contains up to 8 mg of CLO<sub>2</sub> **if fully activated by stomach acid**. The drop size used here is 1/20 of a milliliter or 0.05 ml. This drop size is based on a standard measurement of 20 drops to a milliliter of solution. If you are not sure of your drop size, a 1 ml syringe can be used to measure & dispense MMS drops very accurately. One drop = 0.05 ml, 2 drops = 0.10 ml, 3 drops = 0.15 ml, etc. NOTE: CDS is made with 50% citric acid, not 4% HCL.

The chart below shows the total amount of CLO<sub>2</sub> per hourly dosage if the added MMS is fully activated in stomach acid. For this example, let's assume that you can't ingest more than 5 ml of 3000 ppm CDS hourly, because of mouth and throat irritation.

## Example: Eight Hour Dosing Chart For Adults

(make your own personal chart)

**MMS should be added to each hourly dose and/or to a daily bottle because it is not known how long MMS is effective.**

Dosing Protocol	CDS ml per hour	CDS CLO <sub>2</sub> mg per hour	MMS drops per hour	MMS CLO <sub>2</sub> mg / hour	Total CLO <sub>2</sub> mg / hour	Total CLO <sub>2</sub> mg / 8 doses
A	2	6	-	-	6	48
B	1	3	1	8	11	88
C	2	6	1	8	14	112
D	3	9	1	8	17	136
E	4	12	1	8	20	160
F	3	9	2	16	25	200
G	4	12	2	16	28	224

NOTE: the simple formula for calculating milligrams of CLO<sub>2</sub> is:

**Volume** (liters) x **Concentration** (ppm) = **Dose** (mg of CLO<sub>2</sub>)

*Example:* 1 ml of 3000 ppm CDS = 0.001 x 3000 = 3 mg CLO<sub>2</sub>



3 July 2016

Updated 1Nov'22 and 26Sept'25

<http://mmsinfo.org/>