

➤ If the animal does not show signs of improvement after two days, **move on to Protocol 1000 Plus**. This means add DMSO to each oral dose. Continue using the same dosage amounts for Protocol 1000 and add the following amounts of DMSO:

- Cup1—add 3 drops of DMSO.
- Cup2—add 24 drops of DMSO.
- MMS1 drops—for every 1 drop of MMS1, add 3 drops of DMSO.

Please take special note, that once you add DMSO to the MMS1 mixture in Cup1 and Cup2, it will only maintain full strength for two hours. The "cups" with DMSO added will need to be made up fresh every third hour.

**Column 3: Protocol 2000** dosage for animals. Although Protocol 2000 for people calls for taking MMS1 and MMS2, this column is only for MMS1. This is because normally you would not give an animal MMS2. There are rare exceptions to this however, and I have included details on how to administer MMS2 to animals in column 5. I have only given two numbers in this column. This is because the principle of Protocol 2000 is you work up to taking as many MMS1 drops as you can per hour but without getting sick (in this case without your animal getting sick).

So, the first number is the amount you would begin giving to your animal. This is assuming you have had your animal on Protocol 1000, and worked up to the maximum 3-drop dose equivalent for your animal's weight that Protocol 1000 calls for (as per column 2). At that point, you start increasing the dosage as is called for in Protocol 2000. If you have not worked up to the equivalent 3-drop dose, then start from whatever dosage you are at and